


UNITED STATES NAVAL POSTGRADUATE SCHOOL



SEARCH AND EVASION GAMES

by

Roger G. Schroeder

June, 1966

Technical Report / Research Paper No. 68

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

TA7
.U62
no. 68

UNITED STATES NAVAL POSTGRADUATE SCHOOL
Monterey, California

Rear Admiral E. J. O'Donnell, USN,
Superintendent

Dr. R. F. Rinehart,
Academic Dean

ABSTRACT:

We develop some two-person zero-sum game formulations of search and evasion problems. By employing a game theoretic approach, we allow the hider, as well as the searcher, to choose a strategy. This is in contrast to most search models which assume a stationary or passive hider. Both non-sequential and sequential search games are investigated. Some interesting aspects of the non-sequential game and an example of an antisubmarine search problem are given. The sequential games consist of a sequence of moves. When the players move, they not only determine a payoff but also the probability that the game terminates before the next move. When at most a finite number of moves is allowed, we prove that a solution may be found by solving a recursive sequence of matrix games. When the number of moves is not bounded, the game is characterized by a special type of non-linear program. The solution to this program can be approximated by successive perturbations of a related linear program. Finally, we obtain the result that a pair of strategies minimaxes the expected duration of the game if and only if these strategies also maximin the probability of termination in one step.

Prepared by: R. G. Schröder

Approved by:

J. R. Borsting,

Chairman, Department of
Operations Analysis

Released by:

C. E. Menneken,

Dean of
Research Administration

U. S. Naval Postgraduate School Technical Report / Research Paper No. 68
June, 1966

UNCLASSIFIED

ACKNOWLEDGEMENT

This research was part of the author's Ph. D. dissertation. The dissertation advisor, Dr. A. Charnes of Northwestern University, offered valuable advice and guidance. In addition, this work was undertaken while the author participated in the U.S. Naval Junior Line Officer Advanced Scientific Educational Program.

TABLE OF CONTENTS

	Page
1. Introduction	1
2. The n-cell Search Game	
2.1 Formulation	3
2.2 An Example	5
2.3 Extensions	10
3. The Finite Sequential Game	
3.1 Formulation	11
3.2 Recursive Solution	14
4. The Infinite Sequential Game	
4.1 Formulation	17
4.2 Solution by Perturbation	20
4.3 Iterative Method	25
4.4 Tactical Payoffs and a Special Case	28
References	30

LIST OF FIGURES

Figure	Page
1. The Search Region and Cells	5
2. Probability of Detection versus Range	6
3. Player 1's Payoff Matrix	7
4. Optimal Strategies	9

1. INTRODUCTION

We investigate some two-person zero-sum games which typically arise in search and evasion settings. One of the classical problems in the theory of search is to determine an optimal division of search effort among n -cells.¹ We explore this problem when both the searcher (P1) and the hider (P2) can choose a cell on each move. Here we are allowing for the active participation of the hider as opposed to other formulations which require a stationary hider. An n -cell search game, as we call it, and two sequential extensions of the n -cell game are proposed.

We summarize our game formulations and results. In the n -cell game, every play consists of exactly one move. The payoff function is taken to be the probability that P1 detects P2. This payoff is particularly appealing for Antisubmarine Warfare (ASW) applications. With the indicated payoff, a zero-sum assumption corresponds to the rôle of an evader for P2. To illustrate this point, an example of a typical ASW situation is given. We also show how a constrained-game extension can be employed to include additional tactical information.

The n -cell game is extended to a sequential game. In turn, we consider sequential games consisting of a finite number and an infinite number of moves. On each move, the players not only determine an immediate payoff, but they also determine the probability that the game terminates

¹For an example, see Koopman [9] and Bellman [1].

before the next move. This game may be thought of as a matrix game which is played again with probability determined by the players.

For the finite sequential game, we show how optimal strategies and the value may be computed by dynamic programming. In particular, the solution of the game can be found by solving a recursive sequence of matrix games. The amount of computational effort is usually much less than would be required by solution of the game in normal form.

When an infinite number of moves is allowed, we show how to characterize the resulting sequential game by a special type of non-linear program. If one of the variables in the constraint set of this non-linear program is held fixed, it becomes a linear program. To find the solution of the sequential game, we must adjust this variable in the constraint set to make the optimal value of the objective function equal to zero. We show how to perturb the linear program and thereby approximate the game solution to within desired accuracy.

Many of the search models which appear in the literature assume a stationary hider. Models of this type are given by Koopman [9], Bellman [1], Pollock [15], Dobbie [7], and MacQueen [12]. On the other hand, game formulations which allow the hider to choose strategies are presented by von Neumann [19], Norris [14], and Neuts [13]. Our games are generalizations and extensions of these search games. In particular, one of von Neumann's [19] search games is a special case of the n -cell search game which is presented in section 2.

2. THE N-CELL SEARCH GAME

2.1 Formulation

To formulate the n-cell search game, we assume that the searcher has one detection device and that there is only one hider. Later, we relax these assumptions. The search region of interest is divided into n-cells. A pure strategy for the searcher (P1) is a cell to search (locate his detection device) and a pure strategy for the hider (P2) is a cell in which to hide. A play consists of exactly one simultaneous choice of strategies (move) or, of course, the players may choose their strategies sequentially provided the second choice is made in ignorance of the first.

To define the payoffs, we assume that if P1 looks in cell i and P2 hides in cell j , then P1 detects P2 with probability a_{ij} ($i, j = 1, \dots, n$). Let A be the $n \times n$ payoff matrix $A = (a_{ij})$.

To embrace tactical encounters, we have postulated the payoff as a probability of detection. Other search payoffs could be used as well. We have also allowed the probability of detection to be a function of the range between P1 and P2. This feature is not included in most other detection models.

Now, suppose that P1 searches cell i with probability x_i , and suppose that P2 hides in cell j with probability y_j . We require that

$$\sum_{i=1}^n x_i = 1, \quad x_i \geq 0, \quad i = 1, \dots, n,$$


$$\sum_{j=1}^n y_j = 1, \quad y_j \geq 0, \quad j = 1, \dots, n.$$

Let X and Y be the $n \times 1$ vectors $X = (x_1, \dots, x_n)$ and $Y = (y_1, \dots, y_n)$. Then X and Y are mixed strategies for P_1 and P_2 , respectively. If P_1 chooses X and P_2 chooses Y then, from elementary probability, P_1 detects P_2 with probability $X^t A Y$.¹ We assume that P_1 chooses X to maximize his detection probability $X^t A Y$. Now consider the case when P_2 chooses Y to minimize $X^t A Y$. Then we can interpret the motive of P_2 as evasive action since P_2 is attempting to minimize the probability that he is detected. This action by P_2 also gives rise to a zero-sum game. It is important to note this relationship between evasion and a zero-sum game. We henceforth restrict our discussion to zero-sum games.

¹ X^t denotes the vector X -transpose.

2.2 An Example

To illustrate the n -cell game, consider the following tactical example. Suppose a submarine must pass through a channel to get from its base to its patrol area.¹ The searcher, P_1 , wishes to locate a detection device in the channel to detect submarines as they pass through. For convenience, we assume that the search region is divided into 15 cells, as shown in Figure 1. We assume that P_1 wants to locate his device in one of these cells to maximin the probability of detection.




The Search Region and Cells

Figure 1

¹This type of situation was encountered in the Bay of Biscay during World War II, Sternhell and Thorndike [18].

To obtain the payoff matrix, we need the detection probabilities. These probabilities are found from the probability of detection versus range curve, as shown in Figure 2.



Probability of Detection versus Range

Figure 2

To find the payoff matrix, suppose that P1 locates his detection device in cell 5, and P2 hides in cell 8; then the range is three cells, and the probability of detection, from Figure 2, is $a_{52} = 0.367$. The other elements of the payoff matrix A are determined in a similar manner, and the complete matrix is given in Figure 3.

PLAYER 2'S PURE STRATEGIES

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15

.900 .815 .603 .367 .182 .073

.815 .900 .815 .603 .367 .182 .073

.603 .815 .900 .815 .603 .367 .182 .073

.367 .603 .815 .900 .815 .603 .367 .182 .073

.182 .367 .603 .815 .900 .815 .603 .367 .182 .073

.073 .182 .367 .603 .815 .900 .815 .603 .367 .182 .073

.073 .182 .367 .603 .815 .900 .815 .603 .367 .182 .073

.073 .182 .367 .603 .815 .900 .815 .603 .367 .182 .073

.073 .182 .367 .603 .815 .900 .815 .603 .367 .182 .073

.073 .182 .367 .603 .815 .900 .815 .603 .367 .182 .073

.073 .182 .367 .603 .815 .900 .815 .603 .367 .182

.073 .182 .367 .603 .815 .900 .815 .603 .367

.073 .182 .367 .603 .815 .900 .815 .603

.073 .182 .367 .603 .815 .900 .815

.073 .182 .367 .603 .815 .900

PLAYER 1'S PURE STRATEGIES

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

All blank elements are zeros.

Player 1's Payoff Matrix

Figure 3

The value and optimal strategies (solution) of the game can now be computed. The solution was computed by linear programming, and it is tabulated in Figure 4. From this figure, we observe how the boundaries of the search region affect the searching strategy. A substantial amount of the effort is allocated to the end cells. Notice that the only data required for this search model is a probability of detection versus range curve.

j	y_j
1	0.258
5	0.069
6	0.165
10	0.234
15	0.274

all other $y_j = 0$

j	y_j
1	0.274
6	0.234
10	0.165
11	0.069
15	0.258

all other $y_j = 0$


j	x_j
2	0.317
6	0.099
8	0.168
10	0.099
14	0.317

all other $x_j = 0$

P2's Optimal Basic Strategies


P1's Optimal Strategy

4a



P2's Optimal Strategies

4b



P1's Optimal Strategy

4c

OPTIMAL STRATEGIES

Figure 4

2.3 Extensions

Before leaving this subject, we point out some possible extensions. Additional probabilistic information can be included in the n -cell game by considering a constrained game extension. The elegant development of a constrained game by Charnes [2] can then be applied directly. To illustrate the type of constraints which may arise, suppose that the searcher can bound the probability that the hider chooses certain cells, i. e., the searcher determines numbers L_j and U_j ($0 \leq L_j \leq U_j \leq 1$) such that

$$(1) \quad L_j \leq y_j \leq U_j .$$

These bounds may arise from intelligence or previous contacts.

Constraints of the type in equation (1) and, in general, any linear inequalities can be included in a constrained game formulation. The method of Charnes [2] can then be employed.

It is also desirable to relax the assumption that P1 has only one detection device. This can be easily accomplished by redefining P1's pure strategies in the following way. For simplicity, suppose P1 has two detection devices. Then let each pure strategy for P1 be the two-tuple (k, i) where cell k denotes the location of device 1 and cell i is the location of device 2. Now the probability of detection can be calculated for each such pure strategy, and again an ordinary matrix game is obtained. In a similar way, we can also allow P2 to consist of two or more hiders. Next, we formulate a finite sequential version of the n -cell search game.

3. THE FINITE SEQUENTIAL GAME

3.1 Formulation

First, we discuss the elements of the finite sequential game, and then we proceed with the mathematical formulation. A play of the game consists of, at most, a finite number (N) of moves. On each move, when the game has not terminated, the players are faced with a two-person zero-sum game. In our formulation, we shall use the n -cell game as the two-person zero-sum game for each move. When the players move, they each choose a strategy which determines a zero-sum payoff from player 2 to player 1 and a probability that the game terminates before the next move. We wish to find an optimal strategy for each player which minimizes the expected accumulated payments received by player 1.

The recursive optimization technique which we will propose has also been discussed by other authors. Kuhn [10] (1953) gave his theorem on games of perfect recall which paved the way for further work. Shapley [17] (1953) was the first to point out the recursive character of a generalization of this game, although he did not deal with the finite case. Later contributions were made by Bellman [1] (1957), Everett [8] (1957), Zachrisson [22] (1964), and Denardo [6] (1965).

The payoffs and continuation probabilities are now specified. Suppose that P_1 searches cell i and P_2 hides in cell j on move r . Then the payoff

from P2 to P1 is

$$a_{ij}(r) \quad i, j = 1, \dots, n \\ r = 1, \dots, N .$$

Also, when P1 searches cell i and P2 hides in cell j on move r, the game continues until move $r + 1$ with probability

$$p_{ij}(r) \quad i, j = 1, \dots, n \\ r = 1, \dots, N - 1 .$$

We let A_r be the $n \times n$ matrix $A_r = (a_{ij}(r))$ and P_r the $n \times n$ matrix $P_r = (p_{ij}(r))$. Hence, A_r is P1's payoff matrix for move r and P_r is the matrix of continuation probabilities for move r. We assume that the game is zero sum and that P1 is the maximizing player.

Next, we consider strategies for the players. We have assumed that the continuation probability and payoff depend only on the choices available for a particular move. It follows that the game is one of perfect recall as defined by Kuhn [10]. Kuhn's theorem for a game of perfect recall asserts that a "behavior strategy" is optimal. For this particular game, a behavior strategy takes the following form: let X_r and Y_r be mixed strategies over the alternatives available on move r for P1 and P2, respectively. Let $X = (X_1, \dots, X_N)$ be an N-tuple of the above mixed strategies for P1. Then X is a behavior strategy for P1. Similarly, we define a behavior strategy, $Y = (Y_1, \dots, Y_N)$, for P2. Now the following sets of strategies are introduced:

$$X_r = \{x_r\}, \quad Y_r = \{y_r\}, \quad X = \{X\}, \quad Y = \{Y\} .$$

From Kuhn's theorem of perfect recall, the sets \bar{X} and \bar{Y} contain optimal game strategies for P1 and P2, respectively; and we will, therefore, limit our search for optimal strategies to these sets.

The total expected payoff for P1 will be expressed in terms of fixed strategies $X \in \bar{X}$, $Y \in \bar{Y}$ and the given information. If P1 chooses the strategy $X_r \in \bar{X}_r$ for move r and P2 chooses $Y_r \in \bar{Y}_r$, then the payoff to P1 for move r is

$$X_r^t A_r Y_r \quad r = 1, \dots, N$$

and the game continues until move $r + 1$ with probability

$$X_r^t P_r Y_r \quad r = 1, \dots, N - 1.$$

Now the product of the probability that the game continues until move r and the payoff for move r is

$$X_r^t A_r Y_r \prod_{h=1}^{r-1} X_h^t P_h Y_h \quad r = 2, 3, \dots, N.$$

The expected accumulated payoff for N moves, $v_1(X, Y)$, is the sum of the above terms

$$(2) \quad v_1(X, Y) = X_1^t A_1 Y_1 + \sum_{r=2}^N X_r^t A_r Y_r \prod_{h=1}^{r-1} X_h^t P_h Y_h.$$

Since the game has a finite number of moves and a finite number of strategies, it must have a value and optimal strategies.¹ Recall that the sets \bar{X} and \bar{Y} contain optimal strategies. Therefore, the function $v_1(X, Y)$ has at least one saddle point over the sets \bar{X} and \bar{Y} .

¹ von Neumann and Morgenstern [21].

3.2 Recursive Solution

We show how to compute the minimax of equation (2) by a recursive technique. Let \bar{X}_r and \bar{Y}_r denote the sequences of mixed strategies

$$\begin{aligned}\bar{X}_r &= (X_r, X_{r+1}, \dots, X_N) \\ r &= 1, \dots, N \\ \bar{Y}_r &= (Y_r, Y_{r+1}, \dots, Y_N)\end{aligned}$$

Of course, we have $\bar{X}_1 \equiv X$, and $\bar{Y}_1 \equiv Y$. We rewrite equation (2) and also define the scalar functions $v_r(\bar{X}_r, \bar{Y}_r)$ by

$$\begin{aligned}(3) \quad v_r(\bar{X}_r, \bar{Y}_r) &= X_r^t A_r Y_r + (X_r^t P_r Y_r) v_{r+1}(\bar{X}_{r+1}, \bar{Y}_{r+1}) \quad r = 1, \dots, N \\ v_{N+1} &\equiv 0\end{aligned}$$

Now $v_r(\bar{X}_r, \bar{Y}_r)$ may be interpreted as the expected accumulated payments received by P1 on the last $N - r + 1$ moves of the game.

Equation (3) leads us to believe that the recursive optimization technique of dynamic programming can be employed. We will establish this fact by theorem 1. We define $\hat{v}_r, \hat{X}_r, \hat{Y}_r$ by the following equations

$$\begin{aligned}(4) \quad \hat{v}_r &= \max_{X_r \in X_r} \min_{Y_r \in Y_r} \left[X_r^t A_r Y_r + (X_r^t P_r Y_r) \hat{v}_{r+1} \right] \quad r = 1, \dots, N \\ \hat{v}_{N+1} &\equiv 0 \\ &\equiv \hat{X}_r^t A_r \hat{Y}_r + (\hat{X}_r^t P_r \hat{Y}_r) \hat{v}_{r+1}.\end{aligned}$$

The minimax theorem of von Neumann [21] establishes the existence of $\hat{X}_r, \hat{Y}_r, \hat{v}_r$ as defined by equation (4). The following theorem then relates the solutions of equation (4) to the solutions of the sequential game.

Theorem 1: \hat{v}_1 is the value of the sequential game, and

$\hat{X} = (\hat{X}_1, \dots, \hat{X}_N)$, $\hat{Y} = (\hat{Y}_1, \dots, \hat{Y}_N)$ are optimal strategies for P1 and P2, respectively.

Proof: Since $v_1(X, Y)$ from equation (3) is the expected payoff function for the sequential game, a necessary and sufficient condition for \hat{v}_1 to be the value of the game and \hat{X} , \hat{Y} optimal strategies is

$$v_1(X, \hat{Y}) \leq \hat{v}_1 \leq v_1(\hat{X}, Y) \quad \text{all } X \in \mathcal{X} \text{ and } Y \in \mathcal{Y}.$$

We shall show that the above condition is satisfied by \hat{v}_1 , \hat{X} , \hat{Y} as defined by (4). From (4), we have

$$(5) \quad X_r^t A Y_r + (X_r^t P_r Y_r) \hat{v}_{r+1} \leq \hat{v}_r \leq \hat{X}_r^t A Y_r + (\hat{X}_r^t P_r Y_r) \hat{v}_{r+1},$$

$$\text{all } X_r \in \mathcal{X}_r, Y_r \in \mathcal{Y}_r.$$

We begin an inductive argument

$$v_N(\hat{X}_N, Y_N) = \hat{X}_N^t A_N Y_N \geq \hat{v}_N \quad \text{all } Y_N$$

assume

$$v_{r+1}(\hat{X}_{r+1}, \bar{Y}_{r+1}) \geq \hat{v}_{r+1} \quad \text{for some } r \text{ and all } \bar{Y}_r.$$

By definition,

$$v_r(\hat{X}_r, \bar{Y}_r) = \hat{X}_r^t A_r Y_r + (\hat{X}_r^t P_r Y_r) v_{r+1}(\hat{X}_{r+1}, \bar{Y}_{r+1}).$$

By the inductive assumption and $\hat{X}_r^t P_r Y_r \geq 0$,

$$v_r(\hat{X}_r, \bar{Y}_r) \geq \hat{X}_r^t A_r Y_r + (\hat{X}_r^t P_r Y_r) \hat{v}_{r+1} \quad \text{all } \bar{Y}_r.$$

From the preceding equation and equation (5)

$$v_r(\hat{\bar{X}}_r, \bar{Y}_r) \geq \hat{v}_r \quad \text{all } \bar{Y}_r.$$

Hence, by induction on r ¹

$$v_1(\hat{X}, Y) \geq \hat{v}_1 \quad \text{all } Y \in Y.$$

Similarly, we may establish

$$v_1(X, \hat{Y}) \leq \hat{v}_1 \quad \text{all } X \in X;$$

therefore,

$$v_1(X, \hat{Y}) \leq \hat{v}_1 \leq v_1(\hat{X}, Y) \quad \text{all } X \in X, Y \in Y$$

and the theorem is true.

In order to find the value and optimal strategies, we can solve equation (4) recursively. But, each iteration of equation (4) requires the solution of an ordinary matrix game. Now the solutions (value and optimal strategies) of a matrix game can be found by solving a linear programming formulation of the game.² Hence, we can find the solutions of the sequential game by optimizing a sequence of N linear programs. Of course, the amount of computational effort required by this method will usually be substantially less than the amount required to solve the sequential game in direct normal form.

¹ Notice that by definition of \bar{X}_1 we have $\hat{\bar{X}}_1 \equiv \hat{X} \equiv (\hat{X}_1, \hat{X}_2, \dots, \hat{X}_N)$.

² See Charnes and Cooper [3].

4. THE INFINITE SEQUENTIAL GAME

4.1 Formulation

In this section, we allow an infinite number of moves in the sequential game. Before giving an analytic formulation, we discuss some of the features of the game. In the infinite sequential game, we do not assume a maximum number of moves. The continuation probabilities alone control the termination of the game. However, we assume that the same payoff matrix and the same continuation probability matrix are specified for all moves. We further assume that the probability of continuing until the next move is strictly less than one for all pairs of strategies. This assumption guarantees boundedness of the expected accumulated payments received by P_1 ; and it guarantees that the game terminates with probability one, although the number of moves may not be bounded. Now we turn to a formal definition of the game under consideration.

As in the previous games, we assume that a search region is specified and that it is divided into n cells. If P_1 chooses cell i ($i = 1, \dots, n$) and P_2 chooses cell j ($j = 1, \dots, n$) on move r ($r = 1, 2, \dots$), then P_1 receives from P_2 the payoff

$$a_{ij}$$

and the game continues until move $r + 1$ with probability

$$(6) \quad 0 \leq p_{ij} < 1.$$

Let P be the $n \times n$ matrix $P = (p_{ij})$ and A the $n \times n$ matrix $A = (a_{ij})$. A is the payoff matrix, and P is the matrix of continuation probabilities for every move. We further assume that the game is zero sum and that P_1 is the maximizing player.

The game which we have defined above is one of "perfect recall"; and by Kuhn's [10] theorem, a "behavior strategy" is optimal. If a player uses a behavior strategy, he plays the same mixed strategy over the alternatives in an information set each time the information set is reached, regardless of the past history of the game. Since the matrices A and P apply to every move, the game has only one information set. Therefore, a behavior strategy is simply a mixed strategy which is used for every move of the game. We restrict our attention to these strategies.

Let $X = (x_1, \dots, x_n)$ and $Y = (y_1, \dots, y_n)$ be behavior strategies (mixed strategies over the alternatives) for P_1 and P_2 , respectively. For example, P_1 chooses alternative i with probability x_i on every move. The expected accumulated payment received by P_1 , $v(X, Y)$, when P_1 chooses X and P_2 chooses Y , is simply the sum over all r of the probability that the game lasts until move r times the payment to P_1 for move r ,

$$(7) \quad v(X, Y) = \sum_{r=0}^{\infty} (X^t P Y)^r X^t A Y.$$

The above sum converges, since (6) implies $0 \leq X^t P Y < 1$ for all strategies X and Y . For convenience, we define the matrix $Q = (q_{ij})$ with $q_{ij} = 1 - p_{ij}$ all i, j . Then Q is the matrix of positive termination

probabilities. Equation (7) may be written as

$$(8) \quad v(X, Y) = \frac{X^t A Y}{1 - X^t P Y} = \frac{X^t A Y}{X^t Q Y} .$$

von Neumann [20] first established the existence of a unique value v and optimal strategies \hat{X} and \hat{Y} for the form in (8), i. e., there exists a unique real number v and strategies \hat{X} , \hat{Y} such that

$$(9) \quad \frac{X^t A \hat{Y}}{X^t Q \hat{Y}} \leq v \leq \frac{\hat{X}^t A Y}{\hat{X}^t Q Y} \quad \text{all strategies } X, Y .$$

An elementary proof of this fact was subsequently given by Loomis [11], and this result is a special case of Shapley's [17] more general "stochastic game". Neuts [13] formulated and solved a special case of the infinite sequential game. His P matrix was a diagonal matrix and his A matrix also had a special form.

4.2 Solution by Perturbation

There are no known methods for computing a solution to equation (9). In this section, we develop a computational method to approximate a solution to any desired degree to accuracy. The method is based on a linear programming formulation of a matrix game with an unknown parameter in the constraints. We show that this parameter is equal to the value of the game if and only if the optimal objective function of the linear program is zero. The remainder of our discussion is then devoted to a method for approximating the required value of the parameter.

To begin, we establish a lemma which relates the solution of the infinite sequential game to the solution of an ordinary two-person zero-sum game.

Lemma: A necessary and sufficient condition for v to be the value of the infinite sequential game and \hat{X} , \hat{Y} optimal strategies is that the two-person zero-sum game with payoff matrix $A - vQ$ has value zero and optimal strategies \hat{X} , \hat{Y} .

Proof: For the matrix game $A - vQ$ to have value zero and optimal strategies \hat{X} , \hat{Y} , it is necessary and sufficient that

$$(10) \quad X^t (A - vQ) \hat{Y} \leq 0 \leq \hat{X}^t (A - vQ) Y \quad \text{all strategies } X, Y.$$

But, $X^t Q Y > 0$ for all strategies, X, Y . Hence, v, X, Y satisfy (10) if and only if

$$(10a) \quad \frac{X^t A \hat{Y}}{X^t Q \hat{Y}} \leq v \leq \frac{\hat{X}^t A Y}{\hat{X}^t Q Y} \quad \text{all strategies } X, Y.$$

Equation (10a) is a necessary and sufficient condition for v to be the value of the infinite sequential game and \hat{X} , \hat{Y} optimal strategies. Hence, the lemma is true.

This lemma immediately suggests a method for computing v . The general procedure is to choose a number s and compute the value of the game $A - sQ$. If the value of $A - sQ$ is zero, then $s = v$ and we are finished. If the value of $A - sQ$ is not zero, then we want to choose a new value of s , say s_1 , such that the value of $A - s_1Q$ is "closer" to zero than the value of $A - sQ$. We begin by formulating the matrix game $A - sQ$ as a linear program.

Consider the linear program

$$\begin{aligned}
 & \text{Max } u_s \\
 (11) \quad & u_s e^t - X^t (A - sQ) \leq 0 \\
 & X^t e = 1 \\
 & X \geq 0
 \end{aligned}$$

where e is the $n \times 1$ vector of all "ones", X is an $n \times 1$ vector, s is a fixed scalar, and u_s is a scalar variable. Let \hat{u}_s , \hat{X} be an optimal solution to (11). Then from Charnes [2], \hat{X} is an optimal strategy for P1 and \hat{u}_s is the value of the game $A - sQ$, (s fixed). Of course, an optimal strategy \hat{Y} for P2 is part of an optimal solution to the dual of (11), and \hat{Y} is available when (11) is solved by the simplex method.

Next, we examine the variation in \hat{u}_s which results from a change

in s . This will allow us to perturb s in such a way that we move \hat{u}_s closer to zero. We consider a perturbation from s to $s + \xi$ in problem (11), and we want to relate \hat{u}_s to $\hat{u}_{s+\xi}$. We add and subtract the vector $\xi X^t Q$ from the constraints of (11) and obtain the following equivalent linear program

$$\begin{aligned}
 & \text{Max } u_s \\
 (12) \quad & u_s e^t - X^t (A - (s + \xi) Q) - \xi X^t Q \leq 0 \\
 & X^t e = 1 \\
 & X \geq 0 .
 \end{aligned}$$

We seek to obtain a linear programming formulation of the game $A - (s + \xi) Q$ from (12). Hence, we let $\bar{q} = \max_{i,j} q_{ij}$, $\bar{q} = \min_{i,j} q_{ij}$ and then for $\xi > 0$

$$(13) \quad \xi \bar{q} e^t \leq \xi X^t Q \leq \xi \bar{q} e^t \quad \text{all strategies } X .$$

Now consider the following linear program

$$\begin{aligned}
 & \text{Max } u' \\
 (14) \quad & u' e^t - X^t (A - (s + \xi) Q) \leq \xi \bar{q} e^t \\
 & X^t e = 1 \\
 & X \geq 0 .
 \end{aligned}$$

Problem (14) is "less constrained" than (12). Therefore, the respective optimal solutions must satisfy ("hats" on the variables denote optimal values)

$$(15) \quad \hat{u}' \geq \hat{u}_s .$$

Notice that the right-hand side of the constraints in (14) is a constant vector. We bring this vector over to the left-hand side of the constraints and make the change of variable

$$(16) \quad u = u' - \xi \overline{q}$$

to obtain the program

$$(17) \quad \begin{aligned} & \text{Max } (u + \xi \overline{q}) \\ & u e^t - X^t (A - (s + \xi) Q) \leq 0 \\ & X^t e = 1 \\ & X \geq 0 . \end{aligned}$$

But, (17) is the desired linear programming formulation of the game $A - (s + \xi) Q$ except for the additive constant $+ \xi q$ in the objective function. Hence,

$$\hat{u} = \hat{u}_s + \xi$$

and

$$\hat{u}' = \hat{u}_s + \xi + \xi \overline{q} .$$

From (15) and the above equation

$$\hat{u}_s + \xi + \xi \overline{q} \geq \hat{u}_s .$$

By using the left-hand side of (13), we get by a similar argument

$$\hat{u}_s + \xi + \xi \overline{q} \leq \hat{u}_s .$$

Thus, for $\xi > 0$

$$(18) \quad \hat{u}_s - \xi \overline{q} \leq \hat{u}_s + \xi \leq \hat{u}_s - \xi \overline{q}, \quad \xi > 0$$

and for $\xi < 0$ we can derive the relationship

$$(19) \quad \hat{u}_s - \xi \bar{q} \leq \hat{u}_{s+\xi} \leq \hat{u}_s - \xi \bar{\bar{q}}, \quad \xi < 0.$$

Equations (18) and (19) give the desired relationships. We can choose a starting value of s and then subsequently perturb \hat{u}_s toward zero.

To assist in choosing a starting value of s , we propose the following method. Two numbers, m and M ($m \leq M$), are determined such that $\hat{u}_m \geq 0$ and $\hat{u}_M \leq 0$. Then since \hat{u}_s is a continuous function of s ,¹ $\hat{u}_s = 0$ for some s in the range $m \leq s \leq M$. Furthermore, this value of s is unique. We would then choose the initial value of s to satisfy $m \leq s \leq M$. Suppose we take m and M to be

$$(20) \quad m = \min_{i,j} \frac{a_{ij}}{q_{ij}}, \quad M = \max_{i,j} \frac{a_{ij}}{q_{ij}},$$

then

$$mq_{ij} \leq a_{ij}, \quad Mq_{ij} \geq a_{ij} \quad \text{all } i, j.$$

From the constraints of (11), we see that

$$\hat{u}_s = \min_j \sum_{i=1}^n \hat{x}_i (a_{ij} - sq_{ij});$$

thus,

$$\hat{u}_m \geq 0, \quad \hat{u}_M \leq 0.$$

With certain restrictions on the elements a_{ij} , we can derive tighter bounds than m and M ; but, the bounds given here are adequate for most applications.

¹ This fact is clear from the foregoing derivation.

4.3 Iterative Method

There are various methods which can be used in connection with equations (18) and (19) to move from one value of s to another. We propose one method here which will automatically change s and drive \hat{u}_s to zero. We find two numbers, b and ξ , for fixed s which will give

$$(21) \quad -b \leq \hat{u}_s + \xi \leq b.$$

By reference to (18) and (19), we find that

$$(22) \quad \xi = \frac{2\hat{u}_s}{\bar{q} + \bar{q}}, \quad b = \frac{|\hat{u}_s| (\bar{q} - \bar{q})}{\bar{q} + \bar{q}}.$$

The following method for approximating a solution to equation (9) can now be started. For convenience, let \hat{u}_i be the optimal solution to (11) when s_i is the value of s in the constraint set.

Approximation Method

1. Choose a starting value s_0 and calculate u_0 from (11).
2. Given s_i and \hat{u}_i , let

$$s_{i+1} = s_i + \frac{2\hat{u}_i}{\bar{q} + \bar{q}}$$

and calculate \hat{u}_{i+1} from (11).

Let $U = (\hat{u}_1, \hat{u}_2, \dots)$ be the sequence generated by the above method. We now show that this sequence does indeed converge absolutely to zero.

By step 2 and equations (21) and (22), we have

$$(23) \quad |\hat{u}_{i+1}| \leq |\hat{u}_i| \frac{\overline{\overline{q}} - \overline{q}}{\overline{q} + \overline{\overline{q}}}.$$

Let

$$a = \frac{\overline{\overline{q}} - \overline{q}}{\overline{q} + \overline{\overline{q}}};$$

notice that $0 \leq a < 1$, and we also have

$$|\hat{u}_n| \leq a^n |\hat{u}_0|.$$

Since $a^n \rightarrow 0$, we have $|\hat{u}_n| \rightarrow 0$. Because of this convergence, we can approximate the value of the game to any desired accuracy by the above method.

We point out a few interesting features of the proposed method.

From step 2, the "driving force" which moves us from s_i to s_{i+1} is directly proportional to the magnitude of \hat{u}_i . This feature serves to drive \hat{u}_i rapidly toward zero. We also observe that when $\overline{\overline{q}} = \overline{q}$, we will have $\hat{u}_1 = 0$. This is a nice feature since $\overline{\overline{q}} = \overline{q}$ implies that all elements of the Q matrix are equal and, therefore, the game reduces to an ordinary matrix game (see equation (8)).

An alternative to the method which we have proposed here is to use the contraction property of the operator T which is defined as

$$T\alpha = \text{Val}[A - \alpha P]$$

where α is a scalar and $\text{Val}[A - \alpha P]$ denotes the value of the matrix game $A - \alpha P$. Shapley [17] has shown in a more general setting that

T^n_α approaches the value of the infinite stochastic game for arbitrary α , and Charnes and Schroeder [5] have shown how to obtain a desired approximation to the value or optimal strategies by using linear programming methods.

4.4 Tactical Payoffs and a Special Case

For tactical purposes, two particular payoffs are appealing. As in the n -cell game, the payoff for each move may be the probability that P1 detects P2 in that move. Then, in both sequential games, the expected accumulated payment received by P1 will be the probability that P1 detects P2 in the game. The other payoff of interest is obtained by taking all $a_{ij}(r) = 1$. Then P1 always receives a payoff of one unit for each move, and the expected accumulated payment received by P1 is simply the expected number of moves. When each move takes the same length of time, we may, of course, also interpret this payoff as the expected duration of the game. In one of the games in Charnes and Schroeder [5], we show how to incorporate both of these payoffs in the same formulation. In particular, P1 attempts to maximize the probability of detection while constraining the expected number of moves to be no more than a specified number.

Finally, we show that the infinite sequential game reduces to an ordinary matrix game when P1 attempts to minimize the expected number of moves. From the above discussion, we take all $a_{ij} = 1$ for the expected accumulated payment to be the expected number of moves. Then equation (8) becomes

$$v(X, Y) = \frac{1}{X^t Q Y} .$$

For search problems, P1 wants to choose a mixed strategy X to

minimize the expected number of moves. Therefore, we seek to solve the equation

$$(24) \quad v = \min_X \max_Y \frac{1}{X^t Q Y} \equiv \frac{1}{\hat{X}^t Q \hat{Y}} .$$

Clearly, v , \hat{X} , and \hat{Y} satisfy (24) if they satisfy

$$\frac{1}{v} = \max_X \min_Y X^t Q Y \equiv \hat{X}^t Q \hat{Y} .$$

Hence, P1 can minimax the expected number of moves by maximizing the probability that the game terminates in one move. This is a noteworthy feature of the infinite sequential game. Of course, in this case our perturbation technique is not required since we can solve the game directly by means of a single linear program.

REFERENCES

1. Bellman, R. Dynamic Programming, Princeton University Press, Princeton, New Jersey, 1957.
- + 2. Charnes, A. "Constrained Games and Linear Programming", Proceedings of the National Academy of Sciences, July, 1953, pp. 639-641.
3. _____ and W. W. Cooper. Management Models and Industrial Applications of Linear Programming, Vols. I and II, Wiley, New York, 1961.
4. _____ and _____. "The Theory of Search: Optimum Distribution of Search Effort", Management Science, Vol. 5, No. 1, October, 1958, pp. 44-50.
5. _____ and R. G. Schroeder. "On Some Stochastic Tactical Antisubmarine Games", Systems Research Memo No. 131, Northwestern University, Evanston, Illinois, August, 1965.
6. Denardo, E. V. "Sequential Decision Processes", Ph. D. Thesis, Northwestern University, Evanston, Illinois, June, 1965.
- + 7. Dobbie, James M. "Search Theory: A Sequential Approach", Naval Research Logistics Quarterly, Vol. 10, No. 4, December, 1963, pp. 323-334.
8. Everett, H. "Recursive Games", Contributions to the Theory of Games, Vol. III, Annals of Mathematics Study No. 39, Princeton University Press, Princeton, New Jersey, 1957.
9. Koopman, B. O. Search and Screening, OEG Report No. 56, Washington, D. C., 1946.
- + 10. Kuhn, H. W. "Extensive Games and the Problems of Information", Contributions to the Theory of Games, Vol. II, Annals of Mathematics Study No. 28, Princeton University Press, Princeton, New Jersey, 1953, pp. 193-216.

11. Lommis, L. H. "On a Theorem of von Neumann", Proceedings of the National Academy of Science, USA, Vol. 32, No. 8, August, 1946, pp. 213-215.
12. MacQueen, James B. "Optimal Policies for a Class of Search and Evaluation Problem", Management Science, Vol. 10, No. 4, July, 1964.
13. Neuts, Marcel F. "A Multistage Search Game", Journal of the Society for Industrial and Applied Mathematics, Vol. 11, No. 2, June, 1963, pp. 502-507.
14. Norris, Richard C. "Studies in Search for a Conscious Evader", M. I. T. Lincoln Lab. Technical Report No. 279, 1962.
15. Pollock, S. M. "Sequential Search and Detection", Technical Report No. 5, Research on the Application of Probabilistic Models to Naval Problems, M. I. T., May, 1964.
16. Schroeder, R. G. "Mathematical Models of Antisubmarine Tactics", Ph. D. Dissertation, Northwestern University, Evanston, Illinois, June, 1966.
17. Shapley, L. S. "Stochastic Games", Proceedings of the National Academy of Science, USA, Vol. 39, 1953, pp. 1095-1100.
18. Sternhell, C. M. and A. M. Thorndike. Antisubmarine Warfare in World War II, OEG Report No. 51, Washington, D. C., 1946.
19. von Neumann, John. "A Certain Two-Person Zero-Sum Game Equivalent to the Optimal Assignment Problem", Contributions to the Theory of Games, Vol. II, Annals of Mathematics Study No. 28, Princeton University Press, Princeton, N. J., 1953.
20. _____. "Über ein Ökonomisches Gleichungssystem und eine Verallgemeinerung des Brouwerschen Fixpunktsatzes", Ergebnisse eines Mathematischen Seminars, No. 8, 1938, pp. 73-83, translated as "A Model of General Economic Equilibrium", Review of Economic Studies, Vol. 13, No. 33, 1945-46, pp. 1-9.

21. von Neumann, John and O. Morgenstern. Theory of Games and Economic Behavior, Princeton University Press, Princeton, New Jersey, 1944.
- * 22. Zachrisson, L. E. "Markov Games", Advances in Game Theory, Annals of Mathematical Study No. 52, Princeton University Press, Princeton, New Jersey, 1964, pp. 211-253.

DOCUMENT CONTROL DATA - R&D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) U. S. Naval Postgraduate School Monterey, California		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED	
		2b. GROUP	
3. REPORT TITLE SEARCH AND EVASION GAMES			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Technical Report / Research Paper No. 68			
5. AUTHOR(S) (Last name, first name, initial) Schroeder, Roger G., Lieutenant, junior grade, U. S. Navy			
6. REPORT DATE June, 1966	7a. TOTAL NO. OF PAGES 32	7b. NO. OF REFS 22	
8a. CONTRACT OR GRANT NO.	9a. ORIGINATOR'S REPORT NUMBER(S)		
b. PROJECT NO.			
c.	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)		
d.			
10. AVAILABILITY/LIMITATION NOTICES Distribution of this document is unlimited.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY	
13. ABSTRACT We develop some two-person zero-sum formulations of search and evasion problems. By employing a game theoretic approach, we allow the hider, as well as the searcher, to choose a strategy. This is in contrast to most search models which assume a stationary or passive hider. Both non-sequential and sequential search games are investigated. Some interesting aspects of the non-sequential game and an example of an antisubmarine search problem are given. The sequential games consist of a sequence of moves. When the players move, they not only determine a payoff but also the probability that the game terminates before the next move. When at most a finite number of moves is allowed, we prove that a solution may be found by solving a recursive sequence of matrix games. When the number of moves is not bounded, the game is characterized by a special type of non-linear program. The solution to this program can be approximated by successive perturbations of a related linear program. Finally, we obtain the result that a pair of strategies minimaxes the expected duration of the game if and only if these strategies also maximin the probability of termination in one step.			

14.	KEY WORDS	LINK A		LINK B		LINK C	
		ROLE	WT	ROLE	WT	ROLE	WT
	Game Theory						
	Search Theory						
	Detection						
	Sequential Game						
	Mathematical Search Model						
	Stochastic Game						

INSTRUCTIONS

1. ORIGINATING ACTIVITY: Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. REPORT SECURITY CLASSIFICATION: Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. GROUP: Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. REPORT TITLE: Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. DESCRIPTIVE NOTES: If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. AUTHOR(S): Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. REPORT DATE: Enter the date of the report as day, month, year; or month, year. If more than one date appears on the report, use date of publication.

7a. TOTAL NUMBER OF PAGES: The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. NUMBER OF REFERENCES: Enter the total number of references cited in the report.

8a. CONTRACT OR GRANT NUMBER: If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. PROJECT NUMBER: Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. ORIGINATOR'S REPORT NUMBER(S): Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. OTHER REPORT NUMBER(S): If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. AVAILABILITY/LIMITATION NOTICES: Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. SUPPLEMENTARY NOTES: Use for additional explanatory notes.

12. SPONSORING MILITARY ACTIVITY: Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. ABSTRACT: Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. KEY WORDS: Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, roles, and weights is optional.

DISTRIBUTION LIST

Documents Department
General Library
University of California
Berkeley, California 94720

Lockheed-California Company
Central Library
Dept. 77-14, Bldg. 170, Flt. B-1
Burbank, California 91503

Naval Ordnance Test Station
China Lake, California
Attn: Technical Library

Serials Dept., Library
University of California, San Diego
La Jolla, California 92038

Aircraft Division
Douglas Aircraft Company, Inc.
3855 Lakewood Boulevard
Long Beach, California 90801
Attn: Technical Library

Librarian
Government Publications Room
University of California
Los Angeles, California 90024

Librarian
Numerical Analysis Research
University of California
405 Hilgard Avenue
Los Angeles, California 90024

Chief Scientist
Office of Naval Research
Branch Office
1030 East Green Street
Pasadena, California 91101

Commanding Officer and Director
U. S. Navy Electronics Lab. (Library)
San Diego, California 92152

General Dynamics/Convair
P.O. Box 1950
San Diego, California 92112
Attn: Engineering Library
Mail Zone 6-157

Ryan Aeronautical Company
Attn: Technical Information
Services
Lindbergh Field
San Diego, California 92112

General Electric Company
Technical Information Center
P.O. Drawer QQ
Santa Barbara, California 93102

Library
Boulder Laboratories
National Bureau of Standards
Boulder, Colorado 80302

Government Documents Division
University of Colorado Libraries
Boulder, Colorado 80304

The Library
United Aircraft Corporation
400 Main Street
East Hartford, Connecticut 06108

Documents Division
Yale University Library
New Haven, Connecticut 06520

Librarian
Bureau of Naval Weapons
Washington, D. C. 20360

George Washington University Library
2023 G Street, N. W.
Washington, D. C. 20006

National Bureau of Standards Library
Room 301, Northwest Building
Washington, D. C. 20234

Director
Naval Research Laboratory
Washington, D. C. 20390
Attn: Code 2027

University of Chicago Library
Serial Records Department
Chicago, Illinois 60637

Documents Department
Northwestern University Library
Evanston, Illinois 60201

The Technological Institute, Library
Northwestern University
Evanston, Illinois 60201

Librarian
Purdue University
Lafayette, Indiana 47907

Johns Hopkins University Library
Baltimore
Maryland 21218

Martin Company
Science-Technology Library
Mail 398
Baltimore, Maryland 21203

Scientific and Technical Information
Facility
Attn: NASA Representative
P.O. Box 5790
Bethesda, Maryland 20014

Documents Office
University of Maryland Library
College Park, Maryland 20742

The Johns Hopkins University
Applied Physics Laboratory
Silver Spring, Maryland
Attn: Document Librarian

Librarian
Technical Library, Code 245L
Building 39/3
Boston Naval Shipyard
Boston, Massachusetts 02129

Massachusetts Institute of Technology
Serials and Documents
Hayden Library
Cambridge, Massachusetts 02139

Technical Report Collection
303A, Pierce Hall
Harvard University
Cambridge, Massachusetts 02138
Attn: Mr. John A. Harrison, Librarian

Alumni Memorial Library
Lowell Technological Institute
Lowell, Massachusetts

Librarian
University of Michigan
Ann Arbor, Michigan 48104

Gifts and Exchange Division
Walter Library
University of Minnesota
Minneapolis, Minnesota 55455

Reference Department
John M. Olin Library
Washington University
6600 Millbrook Boulevard
St. Louis, Missouri 63130

Librarian
Forrestal Research Center
Princeton University
Princeton, New Jersey 08540

U. S. Naval Air Turbine Test Station
Attn: Foundational Research Coordinator
Trenton, New Jersey 08607

Engineering Library
Plant 25
Grumman Aircraft Engineering Corp.
Bethpage, L. I., New York 11714

Librarian
Fordham University
Bronx, New York 10458

U. S. Naval Applied Science Laboratory
Technical Library
Building 291, Code 9832
Naval Base
Brooklyn, New York 11251

Librarian
Cornell Aeronautical Laboratory
4455 Genesee Street
Buffalo, New York 14225

Central Serial Record Dept.
Cornell University Library
Ithaca, New York 14850

Columbia University Libraries
Documents Acquisitions
535 W. 114 Street
New York, New York 10027

Engineering Societies Library
345 East 47th Street
New York, New York 10017

Library-Serials Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Librarian
Documents Division
Duke University
Durham, North Carolina 27706

Ohio State University Libraries
Serial Division
1858 Neil Avenue
Columbus, Ohio 43210

Commander
Philadelphia Naval Shipyard
Philadelphia, Pennsylvania 19112
Attn: Librarian, Code 249c

Steam Engineering Library
Westinghouse Electric Corporation
Lester Branch Postoffice
Philadelphia, Pennsylvania 19113

Hunt Library
Carnegie Institute of Technology
Pittsburgh, Pennsylvania 15213

Documents Division
Brown University Library
Providence, Rhode Island 02912

Central Research Library
Oak Ridge National Laboratory
Post Office Box X
Oak Ridge, Tennessee 37831

Documents Division
The Library
Texas A & M University
College Station, Texas 77843

Librarian
LTV Vought Aeronautics Division
P.O. Box 5907
Dallas, Texas 75222

Gifts and Exchange Section
Periodicals Department
University of Utah Libraries
Salt Lake City, Utah 84112

Defense Documentation Center (DDC)
Cameron Station
Alexandria, Virginia 22314
Attn: IRS (20 copies)

FOREIGN COUNTRIES

Engineering Library
Hawker Siddeley Engineering
Box 6001
Toronto International Airport
Ontario, Canada
Attn: Mrs. M. News, Librarian

Exchange Section
National Lending Library for
Science and Technology
Boston Spa
Yorkshire, England

The Librarian
Patent Office Library
25 Southampton Buildings
Chancery Lane
London W. C. 2., England

Librarian
National Inst. of Oceanography
Wormley, Godalming
Surrey, England

Dr. H. Tiderschiold, Director
Library
Chalmers University of Technology
Gibraltargatan 5
Gothenburg S, Sweden

TA7
.U62 Schroeder
no.68 Search and evasion
games.

88199

15 NOV 66
9 DEC 66
29 DEC 66

30 APR 68

12 SEP 68

26 JUL 71

16 OCT 73

DISPLA
8888
14331
16720
16907
18911
18919
21009

TA7
.U62 Schroeder
no.68 Search and evasion
games.

88199

genTA 7.U62 no.68

Search and evasion games.



3 2768 001 61421 7

DUDLEY KNOX LIBRARY